



---

Year: 2020

---

**Constrictores Oppel, 1811 – the available name for the taxonomic group  
uniting boas and pythons**

Georgalis, Georgios L ; Smith, Krister T

**Abstract:** Recent advances in the phylogenetic relationships of snakes using both molecular and morphological data have generally demonstrated a close relationship between boas and pythons but also induced nomenclatural changes that rob the least inclusive clade to which both belong of a name. This name would be tremendously useful, because it is the least inclusive group to which a large number of fossil boa-like or python-like taxa can be assigned. Accordingly, an update of higher-level nomenclature is desirable. We herein provide an overview of all the names that have historically been applied to boas and pythons. We show that the earliest name for the supra-familial group encompassing boas and pythons is Constrictores Oppel, 1811. We herein revalidate it as an order-group name below Alethinophidia Nopcsa, 1923 and provide a phylogenetic definition of it to encompass the modern concepts of Booidea and Pythonoidea. We provide emended diagnoses for Constrictores, Booidea, and Pythonoidea based on recent morphological data-sets. Key words Booidea, nomenclature, Pythonoidea, Serpentes.

DOI: <https://doi.org/10.26049/VZ70-3-2020-03>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-191269>

Journal Article

Published Version

Originally published at:

Georgalis, Georgios L; Smith, Krister T (2020). Constrictores Oppel, 1811 – the available name for the taxonomic group uniting boas and pythons. *Vertebrate Zoology*, 70(3):291-304.

DOI: <https://doi.org/10.26049/VZ70-3-2020-03>

# Constrictores Oppel, 1811 – the available name for the taxonomic group uniting boas and pythons

GEORGIOS L. GEORGALIS<sup>1,2,\*</sup> & KRISTER T. SMITH<sup>3,4</sup>

<sup>1</sup>Dipartimento di Scienze della Terra, Università di Torino, Via Valperga Caluso 35, Torino, 10125, Italy — <sup>2</sup> Department of Ecology, Laboratory of Evolutionary Biology, Faculty of Natural Sciences, Comenius University in Bratislava, Mlynská dolina, Bratislava, 84215, Slovakia —

<sup>3</sup> Department of Messel Research and Mammalogy, Senckenberg Research Institute and Natural History Museum, Senckenberganlage 25, Frankfurt am Main, 60325, Germany — <sup>4</sup> Faculty of Biological Sciences, Institute for Ecology, Diversity and Evolution, Max-von-Laue-Straße 13, University of Frankfurt, 60438 Frankfurt am Main, Germany — \* Corresponding author; dimetrodon82@gmail.com

Submitted April 25, 2020.

Accepted June 17, 2020.

Published online at [www.senckenberg.de/vertebrate-zoology](http://www.senckenberg.de/vertebrate-zoology) on June 26, 2020.

Published in print Q3/2020.

Editor in charge: Uwe Fritz

## Abstract

Recent advances in the phylogenetic relationships of snakes using both molecular and morphological data have generally demonstrated a close relationship between boas and pythons but also induced nomenclatural changes that rob the least inclusive clade to which both belong of a name. This name would be tremendously useful, because it is the least inclusive group to which a large number of fossil boa-like or python-like taxa can be assigned. Accordingly, an update of higher-level nomenclature is desirable. We herein provide an overview of all the names that have historically been applied to boas and pythons. We show that the earliest name for the supra-familial group encompassing boas and pythons is *Constrictores* Oppel, 1811. We herein revalidate it as an order-group name below *Alethinophidia* Nopcsa, 1923 and provide a phylogenetic definition of it to encompass the modern concepts of *Booidea* and *Pythonoidea*. We provide emended diagnoses for *Constrictores*, *Booidea*, and *Pythonoidea* based on recent morphological data-sets.

## Key words

*Booidea*, nomenclature, *Pythonoidea*, *Serpentes*.

## Introduction

Boas and pythons are among the most well known snakes. They comprise the largest forms, both extinct and extant (MURPHY & HENDERSON, 1997; HEAD *et al.*, 2009), and have fascinated humans at least since Antiquity (SCHNEIDER, 1821). They have a broad geographic distribution, covering almost all continents except Antarctica, being found even in remote oceanic islands (WALLACH *et al.*, 2014). Although many species are widely distributed, the range of other species is rather confined geographically. Especially the latter species are increasingly in need of conservation efforts (e.g., REYNOLDS & HENDERSON, 2018); one of them, *Bolyeria multocarinata* (Boié in Boié, 1827) went extinct in historical times, as recently as the 1980s (DAY, 1989; WALLACH *et al.*, 2014). They have an extensive fossil record, dominating snake as-

semblages throughout the Paleogene in North America and Europe, and achieved a considerable species richness and diversity, with a variety of sizes and habits (RAGE, 1984; SZYNDLAR, 1991; HEAD *et al.*, 2009; GEORGALIS & SCHEYER, 2019). Their fossil record comprises primarily vertebrae and, more rarely, isolated skull bones (e.g., GILMORE, 1938; RAGE, 1984; HOLMAN, 2000; SZYNDLAR & RAGE, 2003; SMITH, 2013; GEORGALIS & SCHEYER, 2019), although complete, articulated skeletons have been found in certain *Konservat-Lagerstätten*, and even rare mummified specimens with scales have been recovered (FILHOL, 1877; ROCHEBRUNE, 1884; SMITH & SCANFERLA, 2016; SMITH *et al.*, 2018; SCANFERLA & SMITH, 2020).

Recent advances in snake systematics based on DNA sequence data, coupled with novel interpretations of

morphology and skeletal anatomy, have lead to the recognition of boas and pythons as distinct superfamily-level taxa (e.g., VIDAL *et al.*, 2007; VIDAL & HEDGES, 2009; BURBRINK & CROTHER, 2011; REYNOLDS *et al.*, 2014). Divergence dates among boas and pythons are estimated to have occurred as early as the Paleocene or the Late Cretaceous (HEAD, 2015; HSIANG *et al.*, 2015; ZHENG & WIENS, 2016). The content of Booidea as a superfamily has been restricted (PYRON *et al.*, 2014) to Neotropical boas (Boidae), American dwarf boas (Ungaliophiinae), the Rainbow Boa and rubber boas (Charininae), the Calabar Burrowing Python (*Calabaria reinhardtii*), the Malagasy boas (Sanziniidae), the Old World sand boas (*Eryx* [including *Gongylophis*]), and the Pacific Island boas (*Candoia*). Pythonoidea, in turn, encompasses Old World pythons (Pythonidae), sunbeam snakes (*Xenopeltis*) and the Mexican Burrowing Python (*Loxocemus bicolor*). Boa- and python-like fossils, the majority of which are known exclusively from vertebrae, frequently cannot be assigned with confidence to the boa or python clades, and a term is now lacking for that higher-level clade (BURBRINK *et al.*, 2020) comprising both (e.g., GEORGALIS & SCHEYER, 2019). The question thus arises, what should be the name for the clade uniting boas and pythons? We demonstrate that there is in fact an available name for this group and we accordingly revalidate and redefine it here.

## Taxonomic history

Several available names in the old literature have been established to denote the taxonomic entity comprising boas and pythons. Many early workers simply assigned all boas to the genus *Boa* Linnaeus, 1758 (e.g., LINNAEUS, 1758; BODDAERT, 1783; GMELIN, 1789; SEETZEN, 1796; LINK, 1807); pythons were instead referred to “couleuvres” (e.g., LACÉPÈDE, 1789) or even formally to the genus *Coluber* Linnaeus, 1758 (e.g., LINNAEUS, 1758; GMELIN, 1789; BONNATERRE, 1790; SHAW, 1802). Others assigned the then-known species only to the genera *Boa*, *Python* Daudin, 1803b, and *Eryx* Daudin, 1803d, but did not provide any higher group name (e.g., DAUDIN, 1803b, d). Schneider (1801) placed several boas and pythons in the genus *Boa*, but still kept *Python molurus* (Linnaeus, 1758) in *Coluber*, and he treated “erycine” species as members of *Anguis* Linnaeus, 1758 (note that in this paper we use the term “erycine” in quotes, because molecular and combined phylogenetic analyses have universally held the taxon Erycinae sensu RAGE [1984] to be polyphyletic). In a similar vein, LATREILLE (1804), although he distinguished boas and pythons, included both in Colubrini, along with the majority of snake genera. DUMÉRIL (1805) used the (apparently informal) name “boas” (with lower-case “b”) to include *Boa*, *Coralle* (i.e., *Corallus* Daudin, 1803b), and *Python*, but also the homalopsid *Huria* Daudin, 1803b (misspelled as *Hurriah*; currently *Cerberus* Cuvier, 1829) and the elapid *Acanthophis*

Daudin, 1803b. DUMÉRIL (1805), however, did not place *Eryx* (misspelled by him as *Erix*) into “boas” but rather treated it as distinct. The same author, nevertheless, included “boas” and *Eryx* in a higher, formal, group, which he named Hétérodermes, that also included a number of other snake groups (DUMÉRIL, 1805).

OPPEL (1811a, b) applied the name Constrictores, in two different publications during the same year, to encompass only the genera *Boa* and *Eryx*. Although the genera *Python* Daudin, 1803b, and *Eryx* Daudin, 1803d, had been already established prior to OPPEL’s (1811a, b) works, the latter author mentioned *Eryx* but not *Python*. In OPPEL’s (1811a:383; 1811b:58) concept of *Boa*, however, besides the mentioned species in his text, he also inserted “etc.”, thus making clear that he did not intend to list all known taxa. Notably, he also listed the species *Boa regia* Shaw, 1802 (currently *Python regius*; WALLACH *et al.*, 2014). It thus appears that OPPEL (1811a, b) intended his Constrictores to encompass all large taxa recognized as booids and pythonoids today; furthermore, he specifically excluded aniliids (which he placed in *Tortrix* Oppel, 1811b) from his Constrictores. The name Constrictores was subsequently used by FISCHER VON WALDHEIM (1813), who, however, expanded the denoted group to include the hydrophiine elapids *Platurus* Latreille in Sonnini and Latreille, 1801 (i.e., *Laticauda* Laurenti, 1768) and *Hydrophis* Latreille in Sonnini and Latreille, 1801.

RAFINESQUE (1815) applied the name Aplepia to encompass *Boa*, *Python*, *Eryx*, and *Corallus*, as well as the peculiar homalopsid genus *Erpeton* Lacépède, 1801. RAFINESQUE (1815:77) united these genera into Aplepia on the basis of their shared presence of “[u]n seul rang de plaques sous la queue ou le corps” [a single row of plates underneath the body]; according to his taxonomic scheme, Aplepia was the counterpart of Diplepia (including *Coluber* and certain other colubrids, but also the elapid *Acanthophis*) and both of them together constituted the family Colubrinia, which in turn belonged to the suborder Heterodermia.

CUVIER (1817) placed boas in their own group, termed Boas, subdivided into “Boas propres”, *Erix* (sic), and Erpetons (including *Erpeton*), whereas he placed pythons (his Pythons) along with colubrids in a different group, Couleuvres. The same author subsequently kept practically the same taxonomic arrangement, with his Boas being distinct from his Pythons, the latter still pertaining to Couleuvres (CUVIER, 1829).

MAYER (1824) created the name Phaenopoda, denoting the presence of “spur-like” hind-limb rudiments in these snakes, to encompass *Boa*, *Python*, and *Eryx*, as well as *Tortrix* (i.e., aniliids). That name was later discussed by DUMÉRIL & BIBRON (1844; as “Phénopodes”), but curiously the name Phaenopoda was not mentioned at all in the detailed catalog of BOULENGER (1893).

GRAY (1825) established Boidæ to include both boas and pythons, an arrangement that the same author continued to follow (e.g., GRAY, 1842, 1849, 1858). This arrangement met wide acceptance among prominent ophidian researchers from the second half of the 19th through

the 20th centuries, who all placed these snakes together into a single family, Boidae (e.g., CANTOR, 1847; Boæidæ of JAN, 1863; BOULENGER, 1890, 1893, 1913; SCLATER, 1891; ZACHARIAS, 1898; ZENNECK, 1898). In this scheme, until recently, boas and pythons were usually distinguished at the subfamily level, as Boinae and Pythoninae respectively (GADOW, 1909; FRASER, 1937; HOFFSTETTER, 1939, 1955; KUHN, 1946, 1963; ROMER, 1956; FRAZETTA, 1959, 1966; ROUX-ESTÈVE, 1965; RAGE, 1984; UNDERWOOD & STIMSON, 1990; SZYNDLAR, 1991; HOLMAN, 2000; IVANOV, 2000; SZYNDLAR & RAGE, 2003).

HAWORTH (1825) placed *Boa*, *Eryx*, and *Python*, as well as a number of other non-venomous snakes in a group termed *Innocua*, this being the counterpart of all venomous snakes, which were forming the group *Venennata*. A similar arrangement of snakes on the basis of venomousness continued to appear in the 19th century literature (e.g., *Serpenti Innocui* and *Serpenti Velenosi* of FILIPPI, 1840).

FITZINGER (1826) also placed both boas and pythons in a single group, which he called *Pythonoidea*. He later included them in *Saurophidia*, which comprised *Gongylophes* (*Eryx* and allied forms), *Centrophes* (boas), and *Pythophes* (pythons), along with scolecophidians and aniliids (FITZINGER, 1843). Still later, he emended the spelling of *Pythophes* to *Pythonophes* (FITZINGER, 1861). The names *Centrophes* (for boas) and *Pythophes* (for pythons) were subsequently adopted by DIESING (1851), who in his taxonomic scheme, however, omitted *Eryx*. The name *Pythonoidea* for both boas and pythons was used by LEUCKART (1841), whereas EICHWALD (1831) and SCHINZ (1833–1835) used the spellings *Pythonoidei* and *Pythonoideae* respectively for the same group.

RITGEN (1826) used the name *Onychophori* to denote the group encompassing the genera *Eryx*, *Python*, and *Boa*, but also *Tortrix* (i.e., aniliids).

Curiously, Boié (1826, 1827) placed *Boa*, *Python*, and *Eryx* into *Colubrini*. BONAPARTE (1831) used the name *Boidæ* to encompass the subgroups *Boina* (including both *Boa* and *Python*) and *Erycina* but also *Typhlopodina* (i.e., scolecophidians). The same author subsequently modified his taxonomic scheme, removed scolecophidians from snakes, but still treated *Eryx* as distinct from boas and pythons, thus recognizing the families *Erycidae* (including *Erycina* but also *Calamarina* [i.e., the caenophidian *Calamaria* Boié H. in Boié F., 1827]) and *Boidæ* (including *Boina* and *Pythonina*).

MÜLLER (1831) used the name *Macrostomata* to differentiate large-gaped snakes from their counterpart *Microstomata* (which included scolecophidians, aniliids, but also amphisbaenians). The concept of *Macrostomata* includes, in addition to boas and pythons, an array of other snake taxa; indeed, under modern phylogenetic concepts, this taxonomic entity is now used to encompass all alethiophidians to the exclusion of *Anilius*, cylindrophiids, and uropeltids (e.g., LEE & SCANLON, 2002; WILCOX *et al.*, 2002; HSIANG *et al.*, 2015; but see Burbrink *et al.*, 2020). Note also, however, that contrary to the widespread misconception that MÜLLER (1831) created that name, Mac-

rostomata was first used by RITGEN (1826). Within *Macrostomata* MÜLLER (1831) distinguished *Holodonta* (for pythons) and *Isodontia* or *Aproterodonta* (for boas).

The name *Peropodes* appeared first in WIEGMANN (1832) to denote the group encompassing all boas, erycines, and pythons and quickly became among the most widely used in the 19th century literature (FEDOROVICH GORIANINOW, 1834; BURMEISTER, 1837; SCHUBERT, 1837; GORSKI, 1852; TROSCHEL, 1861; MEYER, 1874; SCHREIBER, 1875; MÜLLER, 1878, 1880; PETERS, 1882; HOFFMAN, 1890; GIRARD, 1895). It was later emended as *Peropodae* (BREHM, 1878; BEDRIAGA, 1882) or *Peropoda* (LICHENSTEIN & MERTENS, 1856; COPE, 1862, 1886, 1893, 1894, 1895, 1898), and this name was even, although sporadically, also used during the 20th century (e.g., KRITZESCU, 1902; NOGUCHI, 1909; LEBLANC, 1920; GILMORE, 1938). Indeed, the name *Peropodes* seems to have been so widespread that even FITZINGER (1867) adopted it over names he had previously used for the group encompassing boas and pythons as well as *Uropeltis* Cuvier, 1829, *Xenopeltis* Reinwardt in Boié, 1827, and (the currently much distantly related) *Calamaria*.

SCHLEGEL (1837) included a group termed *Boas* the genera *Boa* and *Python* as well as the caenophidian *Acrochordus* Hornstedt, 1787.

SWAINSON (1839) placed *Boa*, *Python*, and *Eryx* in *Coluberidæ*, along with a large array of other, non-venomous, snake genera, mainly *Colubriformes* (sensu ZAHER *et al.*, 2009).

In their monumental work, DUMÉRIL & BIBRON (1844) introduced the name *Azémiophides* to encompass all boas (*Boæides* and *Érycides*), pythons (*Pythonides*), and *ungaliophiines* as well as aniliids (*Tortricides*). A few years later, they redefined the concept of *Azémiophides*, proposing also the alternative name *Aglyphodontes*, which encompassed the aforementioned taxa plus acrochordids, uropeltids, and certain *Colubriformes* (DUMÉRIL, 1853; DUMÉRIL *et al.*, 1854a, b). They still, nevertheless, treated boas and erycines as a group (*Aprotérodontes*) distinct from that of pythons (*Holodontes* or *Holodontiens*) (DUMÉRIL & BIBRON, 1844; DUMÉRIL, 1853, 1859; DUMÉRIL *et al.*, 1854a, b) and this arrangement continued even later (ROCHEBRUNE, 1880).

The name *Asinea* was used by COPE (1864) to denote the group encompassing *Xenopeltis*, pythons, boas as well as *Acrochordus*, which he all, nevertheless, still treated as distinct families (*Xenopeltidæ*, *Pythonidæ*, *Boidæ*, and *Acrochordidæ* respectively). This distinction of *Pythonidae* from *Boidæ* was subsequently followed by the same author (COPE, 1893) and some other prominent workers (GÜNTHER, 1864; ZITTEL, 1887–1890; LYDEKKER, 1888; HOFFMANN, 1890). The name *Asinea* was subsequently also used by GILMORE (1938).

JAN (1865) called the group as *Boidiens*, and further divided it into *Erycides*, *Boæides*, and *Pythonides*. The same author excluded from that group *Xenopeltis*, which he placed instead with “anilioids” (his *Tortriens*).

The name *Boaeformes* has also appeared in the literature (HEILPRIN, 1907; HAAS, 1952), being also circulated

under the alternative spelling Boaeformia (IHERING, 1911; STROMER, 1912; KUHN, 1946).

Hoffstetter (1939) introduced the term Henophidia for a group uniting boas, pythons, and “anilioids”, and this arrangement was followed more or less consistently in the next decades (e.g., HOFFSTETTER, 1955; ROMER, 1956; UNDERWOOD, 1967; GASC, 1974; RIEPPEL, 1977, 1988; GROOMBRIDGE, 1979; HARDING & HOLMAN, 1981; CUNDALL *et al.*, 1993; REYNOLDS *et al.*, 2014). However, Henophidia has now also been used for the clade comprising both boas, pythons, and caenophidians, i.e., alethinophidians to the exclusion of “anilioids”, and sometimes *Xenopeltis* and *Loxocemus* Cope, 1861 (e.g., BURBRINK & CROTHER, 2011; GAUTHIER *et al.*, 2012; FIGUEROA *et al.*, 2016; PETERMANN & GAUTHIER, 2018).

## Current taxonomies

Recent rank-based taxonomies, relying on molecular and/or morphology-based phylogenies, currently treat Boidae and Pythonidae as distinct families, and a number of smaller groups are also separated as different families (DOWLING *et al.*, 1996; SLOWINSKI & LAWSON, 2002; WILCOX *et al.*, 2002; LAWSON *et al.*, 2004; NOONAN & CHIPINDALE, 2006; VIDAL *et al.*, 2007, 2009; VIDAL & HEDGES, 2009; WIENS *et al.*, 2012; PYRON *et al.*, 2013; REYNOLDS *et al.*, 2014; HSIANG *et al.*, 2015; FIGUEROA *et al.*, 2016; STREICHER & WIENS, 2016; ZHENG & WIENS, 2016; HARRINGTON & REEDER, 2017; BURBRINK *et al.*, 2020). The relatives of Boidae and Pythonidae are united as superfamilies, i.e., Booidea and Pythonoidea (e.g., SCANLON & LEE, 2011; PYRON *et al.*, 2014; WALLACH *et al.*, 2014). In particular, the family “Boidae” sensu PYRON *et al.* (2013) and REYNOLDS *et al.* (2014) has been elevated to the superfamily Booidea, containing the families Boidae, Calabariidae, Candoiidae, Charinidae (comprising Charininae and Ungaliophiinae), Erycidae, and Sanziniidae (PYRON *et al.*, 2014), whereas Pythonoidea is conceived as containing Pythonidae, Loxocemidae, and Xenopeltidae (WALLACH *et al.*, 2014). Bolyeriidae (Round Island boas) and Xenophidiidae (comprising only *Xenophidion*) are thought to be closely related to Booidea or Pythonoidea based on molecular analyses (e.g., WALLACH *et al.*, 2014).

Some recent phylogenetic analyses using DNA sequence or combined data have suggested that Pythonoidea may be more closely related to Uropeltidae and/or Cylindrophiidae than to Booidea (LAWSON *et al.*, 2004; OGUIURA *et al.*, 2010; REYNOLDS *et al.*, 2014; TONINI *et al.*, 2016), or even Booidea more closely related to Cylindrophiidae than Pythonoidea (LI *et al.*, 2020), but Booidea and Pythonoidea are otherwise found to be monophyletic with respect to those and other major clades (e.g., SLOWINSKI & LAWSON, 2002; LEE *et al.*, 2007; PYRON *et al.*, 2013; STREICHER & WIENS, 2016). Most recently, BURBRINK *et al.* (2020), considered 394 loci. In their maximum likelihood tree based on analysis of the

partitioned, concatenated dataset, they found bootstrap support of 99.7% (ultrafast bootstrap approximation) and Shimodaira-Hasegawa approximate likelihood ratio test value of 100% for a clade comprising Pythonoidea, Bolyeriidae, and Booidea as defined above (BURBRINK *et al.*, 2020:Supplementary Data S7); such values comprise unambiguous support in ZAHER *et al.*’s (2019) classification. Fossil-calibrated species-tree methods applied to the same dataset recovered the same topology with less strong support (BURBRINK *et al.*, 2020). Thus, the analyses with the broadest taxon sampling (PYRON *et al.*, 2013) and the most in-depth gene sampling (BURBRINK *et al.*, 2020) have come to the same conclusion.

Furthermore, phylogenetic analyses of morphology have also strongly supported a sister-group relationship between boas and pythons (e.g., LEE & SCANLON, 2002; GAUTHIER *et al.*, 2012; ZAHER & SCANFERLA, 2012; SCANFERLA & SMITH, 2020), although the position of minor lineages (*Xenopeltis*, *Loxocemus*, Bolyeriidae, *Calabaria*) relative to this clade has vacillated.

Thus, in contrast to the taxon name Iguanidae (= Pleurodonta; Torres-Carvajal *et al.*, in press), there has been no single name referring to a group of similar composition throughout the previous century. In addition to repeated alterations to the extant members of Boidae, we also note that some fossil taxa were previously placed in this taxon (SIMPSON, 1933; GILMORE, 1938; HOFFSTETTER, 1955; RAGE, 1984; HOLMAN, 2000), which are now universally accepted to pertain to other groups, such as the extinct Palaeophiidae and Madtsoiidae (RAGE *et al.*, 2003; WALLACH *et al.*, 2014; GEORGALIS *et al.*, 2020).

## Availability and a phylogenetic definition of Constrictores Oppel, 1811

Taking into consideration the survey of literature presented above, it is evident that the oldest available name for the least inclusive group uniting boas and pythons is Constrictores Oppel, 1811a. As mentioned above, OPPEL (1811a, b) published his squamate classification in two different works: the first (OPPEL, 1811a) was published in the 16<sup>th</sup> volume of *Annales du Muséum d’histoire Naturelle* – this volume is dated on its cover page as “1810”, however, it has been subsequently demonstrated by SHERBORN (1914) that the volume was in fact published in early 1811 (pages 328–428 of the volume were published between January and March of that year). The second work (OPPEL, 1811b), which is by far the most popular and remains a key publication for reptile systematics, was a book published in Munich around December 1811 and dealt with all extant reptile and amphibian groups. It is thus clear that OPPEL (1811a) was published before OPPEL (1811b) and therefore, the name Constrictores first appeared in OPPEL (1811a). Be that as it may, the relevant text about Constrictores (and the inclusive genera *Eryx* and *Boa*) in both OPPEL’s (1811a, b) works was almost identical, with only minor wording differences (e.g., “corpus cylindraceum” in

OPPEL [1811a:382] vs. “corpus cylindricum” in OPPEL [1811b:56]). Another difference between OPPEL (1811a) and (1811b) is that in the latter he used the Latin word “Familia” prior to the word Constrictores (OPPEL, 1811b) but he did not use any such denomination or rank in his earlier (1811a) work.

Note that the name Constrictores is a formal Latin name and not a vernacular of German, French or some other modern language. The word “constrictor” is derived from the Latin verb “constringere” (constrict, strangle). It is masculine substantive of the third declension. Accordingly, the nominative plural carries the suffix -es, viz. “constrictores” (Pr. Patrick Smith, pers. comm., 2019). “Constrictor” is not a French word (Robert French Dictionary). Moreover, the name Constrictores does not originate from the genus name *Constrictor*, as OPPEL (1811a, b) does not mention this genus at all (see also Etymology below). Many other formal Latin names appear in the same OPPEL’s (1811a, b) paper (e.g., Squamata [spelled as “Squammata” in the 1811a paper], Testudinata, Saurii, Ophidii, Colubrini, etc). Almost the whole text of both papers is written in Latin, with only the title and few paragraphs being in French in the first paper (OPPEL, 1811a) and only the title and a few pages being in German in the second paper (OPPEL, 1811b). The fact that the word Constrictores is Latin is further supported by the fact that the immediately succeeding words of its “diagnosis” are also in Latin (“Cauda attenuata, rotundata; tela venenifera nulla; calcaria ad anum” [OPPEL, 1811a:377; OPPEL, 1811b:49]).

OPPEL (1811a) did not provide any rank denomination for Constrictores in his earlier work, though he later used the term “Familia” for that grouping (OPPEL, 1811b). In the modern taxonomic scheme we propose for Constrictores that encompasses both boas and pythons, this grouping is not a family-level one. Therefore it does not necessitate an amendment of the name or its ending, as has been the case with family-level names introduced in OPPEL’s (1811a, b) works (such as Viperini, emended to Viperidae, and Colubrini, emended to Colubridae). The fact that the name was not originally proposed as a family group name helps to avoid one potential complication that would ensue. Namely, even though OPPEL (1811a) did not mention the genus *Constrictor* Laurenti, 1768, one could hypothesize that it could be the type genus of the family Constrictoridae. The ICZN (1999: Article 11.7.1.1) dictates: “a family-group name when first published must be a noun in the nominative plural formed from the stem of an available generic name [Art. 29] (indicated either by express reference to the generic name or by inference from its stem); the generic name must be a name then used as valid in the new family-group taxon [Arts. 63, 64] (use of the stem alone in forming the name is accepted as evidence that the author used the generic name as valid in the new family-group taxon unless there is evidence to the contrary)” and later clarifies that “[the family group name must] be clearly used as a scientific name to denote a suprageneric taxon and not merely as a plural noun or adjective referring to the

members of a genus” (ICZN, 1999: Article 11.7.1.2). In summary, whereas OPPEL (1811a) did clearly use Constrictores as a scientific name, it was not as a family-group name, so there would be no reason to emend it to Constrictoridae.

Most importantly, *Constrictor* is a junior synonym of *Boa* Linnaeus, 1758, with the latter genus mentioned by OPPEL (1811a). The ICZN (1999: Article 40.1) dictates that “when the name of a type genus of a nominal family-group taxon is considered to be a junior synonym of the name of another nominal genus, the family-group name is not to be replaced on that account alone”, so Boidae would still have priority over the hypothetical family level Constrictoridae.

We therefore regard Constrictores Oppel, 1811a as a valid name at the supra-familial level (order-group name). From the point of view of hierarchy, Constrictores is ranked below the level of Alethinophidia Nopcsa, 1923, and above the level of the superfamilies Booidea and Pythonoidea.

The fact that the name Constrictores had virtually no usage or mentions during the 19<sup>th</sup> and 20<sup>th</sup> centuries does not invalidate it or render it obsolete, as would certainly be the case with names at the family, genus, or species level: the latter would eventually be rendered nomina obliterata (ICZN, 1999). As such, there is no criterion of prevailing usage in the case of names above the family level, which could possibly lead to the revalidation of the name Peropodes, which appeared extensively in the 19<sup>th</sup> century literature and also occasionally appeared even in 20<sup>th</sup> century. Thus, we resurrect the name Constrictores Oppel, 1811a as an ordinal-group name encompassing Booidea, Pythonoidea and Bolyeroidea.

Furthermore, we define the name phylogenetically following the PhyloCode (CANTINO & DE QUEIROZ, 2014).

**Constrictores** Oppel, 1811a [Georgalis & Smith, this paper], converted clade name

**Registration number.** 309 [[www.phyloregnum.org](http://www.phyloregnum.org)]

**Definition.** The crown clade originating in the last common ancestor of *Boa constrictor* Linnaeus, 1758, and *Python* (originally *Boa regius*) (Shaw, 1802), provided that neither internal specifier is more closely related to any of the following species than to each other: *Typhlops lumbricalis* (Linnaeus, 1758), *Leptotyphlops nigricans* (Schlegel, 1837–1844), *Anomalepis mexicanus* Jan, 1860, *Anilius scytale* (Linnaeus, 1758), *Uropeltis ceylanicus* Cuvier, 1829, *Cylindrophis rufus* (Laurenti, 1768), and *Coluber constrictor* Linnaeus, 1758. Abbreviated definition: *Constrictores* = <  $\nabla$  *Boa constrictor* Linnaeus, 1758 & *Python regius* (Shaw, 1802), provided that neither internal specifier is more closely related to any of the following species than to each other: *Typhlops lumbricalis* (Linnaeus, 1758), *Leptotyphlops nigricans* (Schlegel, 1837–1844), *Anomalepis mexicanus* Jan, 1860, *Anilius scytale* (Linnaeus, 1758), *Uropeltis ceylanicus* Cuvier,

1829, *Cylindrophis rufus* (Laurenti, 1768), and *Coluber constrictor* Linnaeus, 1758.

Note that in this definition we chose to exclude not only the first-named scolecophidian (i.e., the typhlopoid *Typhlops lumbricallis*) but also the types of the eponymous Linnaean families Leptotyphlopidae and Anomalepididae, taking into consideration that many phylogenetic analyses indicate that Scolecophidia is paraphyletic (e.g., ZHENG & WIENS, 2016; HARRINGTON & REEDER, 2017; MIRALLES *et al.*, 2018; BURBRINK *et al.*, 2020; but see STREICHER & WIENS, 2016). Further note that the eponymous type species of Leptotyphlopidae is *Leptotyphlops nigricans* (Schlegel, 1837–1844) and not *Leptotyphlops albifrons*, i.e., *Stenostoma albifrons* Wagler in Spix and Wagler, 1824 (currently *Epictia albifrons*) as stated by LEE *et al.* (2007).

**Etymology.** As was mentioned above, the word “Constrictores” originates from the Latin verb “constringere” (constrict, strangle). The name of the type species of Boidae, *Boa constrictor*, obviously refers to the snake’s method of killing prey (e.g., BOBACK *et al.*, 2015). Note that the genus names *Constrictor* Laurenti, 1768 and *Constrictor* Wagler, 1830, were independently established to accommodate species of *Boa* and *Python* respectively (as was also mentioned above, OPPEL [1811a, b] made no single mention of the genus name *Constrictor* in his works). Neither should *Constrictores* be confused with the type species of Colubridae, *Coluber constrictor* Linnaeus, 1758. Constriction appears to be a widespread killing method across the different lineages within *Constrictores* (e.g., CUNDALL & IRISH, 1986). Of course, the killing of prey by constriction is a widespread habit among members of other snake clades (e.g., HSIANG *et al.*, 2015). We propose to use the informal term “constrictors” to refer to members of the clade *Constrictores*.

**Primary reference phylogeny.** Burbrink *et al.* (2020), fig. S7; this paper, Fig. 1.


**Composition.** Apart from the taxa subsuming the internal specifiers, i.e., Boidae Gray, 1825, sensu PYRON *et al.* (2014) and Pythonidae Fitzinger, 1826, sensu Wallach *et al.* (2014), the taxonomic content of *Constrictores* following the reference phylogeny and most other molecular and combined analyses is clear: *Eryx*, *Candoia*, *Sanziniidae*, *Charinidae*, *Calabaria*, *Loxocemus*, *Xenopeltis*, *Bolyeriidae*, and *Xenophidion*.

On the booid side, *Eryx* was for long time treated as a distinct family, Erycidae (e.g., Érycidae of DUMÉRIL & BIBRON, 1844, and DUMÉRIL *et al.*, 1854c; JAN, 1862; GÜNTHER, 1864; CARUS, 1868; COPE, 1883; BOETTGER, 1884; HOFFMANN, 1890; POČTA, 1905; STROMER, 1910); since the mid-20<sup>th</sup> century, Erycidae was usually treated as a subfamily of Boidae, as Erycinae (e.g., HECHT, 1959; HOFFSTETTER & RAGE, 1972; RAGE, 1977, 1984; SZYNDLAR, 1991; KLUGE, 1993; SZYNDLAR & SCHLEICH, 1994; SZYNDLAR & RAGE, 2003; BASZIO, 2004; SMITH, 2013; WALLACH *et al.*, 2014). Recent taxonomic schemes place *Eryx* in a

distinct family close to Boidae (e.g., PYRON *et al.*, 2014; FIGUEROA *et al.*, 2016; BURBRINK *et al.*, 2020). Other clear members of *Constrictores* on the booid side are Charinidae Gray, 1849 (sensu PYRON *et al.*, 2014) (comprising Charininae Gray, 1849, and Ungaliophiinae McDowell, 1987), *Candoia* Gray, 1842 (for which the monotypic family Candoiidae Pyron, Reynolds, & Burbrink, 2014 was established), *Calabaria* Gray, 1858 (for which the monotypic family Calabariidae Gray, 1858, was established), and Sanziniidae Romer, 1956 (including *Acrantophis* Jan, 1860, and *Sanzinia* Gray, 1849).

On the pythonoid side, *Xenopeltis* Reinwardt in Boié, 1827 (for which the monotypic family Xenopeltidae Bonaparte, 1845, was established) and *Loxocemus* Cope, 1861 (for which the monotypic family Loxocemidae Cope, 1861, was established) are inferred to be successive sister taxa of Pythonidae in molecular analyses (e.g., SLOWINSKI & LAWSON, 2002; PYRON *et al.*, 2013; REYNOLDS *et al.*, 2014; FIGUEROA *et al.*, 2016; STREICHER & WIENS, 2016; ZHENG & WIENS, 2016; HARRINGTON & REEDER, 2017; BURBRINK *et al.*, 2020), which makes them members of *Constrictores*. Note, however, that in many morphology-only analyses these two lineages fall outside of the clade comprising boas, pythons, and caenophidians (e.g., LEE & SCANLON, 2002; HSIANG *et al.*, 2015). More recent studies have concluded that the Asian *Xenophidion* Günther and Manthey, 1995 (for which the monotypic family Xenophidiidae Wallach & Günther, 1998, was established) and/or the Mascarene Bolyeriidae Hoffstetter, 1946, are related to boas (STREICHER & WIENS, 2016; ZHENG & WIENS, 2016; HARRINGTON & REEDER, 2017; BURBRINK *et al.*, 2020), which would also make them members of *Constrictores*; this conclusion is unchanged if they are more closely related to pythons instead (LAWSON *et al.*, 2004; species-tree analysis of BURBRINK *et al.*, 2020). There is considerably molecular evidence that *Xenophidion* and Bolyeriidae are sister-taxa, starting with LAWSON *et al.* (2004), and they share a synapomorphy that is unique among tetrapods: a jointed maxilla. However, *Xenophidion* was not included in the reference phylogeny of Burbrink *et al.* (2020), so further work is desirable to test its membership.

The case of Tropidophiidae (i.e., extant *Tropidophis* Bibron in Ramón de la Sagra, 1838–1843, and *Trachyboa* Peters, 1860) bears elaboration. They were long lumped into boids (e.g., ROMER, 1956; RAGE, 1984; SZYNDLAR & BÖHME, 1996), including also ungaliophiines (e.g., SZYNDLAR & RAGE, 2003). However, formal phylogenetic analyses of morphology (e.g., LEE & SCANLON, 2002; GAUTHIER *et al.*, 2012; ZAHER & SCANFERLA, 2012; HSIANG *et al.*, 2015; SCANFERLA *et al.*, 2016; SCANFERLA & SMITH, 2020) have generally supported the hypothesis of ZAHER (1994), based on external and muscular morphology, that Tropidophiidae is more closely related to Caenophidia than to Ungaliophiinae, and the latter related to boas. Molecular studies, on the other hand, have suggested a radically different topology, with Tropidophiidae being the sister taxon to *Anilius* (WILCOX *et al.*, 2002; LAWSON *et al.*, 2004; GOWER *et al.*, 2005; VIDAL *et al.*, 2007, 2009;



**Fig. 1.** Reference phylogeny (after BURBRINK *et al.*, 2020: Data File S7). Lizard outgroups were removed and snake outgroups to Constrictores were collapsed. Support values are BS/SH and are 100/100 for all ingroup taxa except where noted. The nodes corresponding to Booidea, Pythonoidea and Constrictores are labeled. Note also that the Xenophidiidae is not included herein, although it could pertain as well to Constrictores, as it was not included in the analysis of BURBRINK *et al.* (2020). Abbreviations: SH, the Shimodaira-Hasegawa likelihood ratio; BS, bootstrap. Photograph of *Eryx jaculus* by Ilias Strachinis; photographs of *Eunectes murinus* and *Python sebae* by Alberto Sanchez Vialas. Images of *Acrantophis dumerili*, *Boyeria multocarinata*, *Charina bottae*, *Loxocemus bicolor*, and *Xenopeltis unicolor* reproduced from JAN & SORDELLI (1860–1866), CALABRIA (1858), and GÜNTHER (1877).

BURBRINK & CROTHER, 2011; REYNOLDS *et al.*, 2014; STREICHER & WIENS, 2016; MIRALLES *et al.*, 2018; BURBRINK *et al.*, 2020). The distinctiveness of Tropidophiidae and Ungaliophiinae is further corroborated by their cranial (BOGERT, 1968a) and vertebral anatomy (BOGERT, 1968a, b; SZYNDLAR & RAGE, 2003; SMITH, 2013). In summary, there is neither morphological nor molecular support for the inclusion of Tropidophiidae in Constrictores.

**Synonyms.** Aplezia of RAFINESQUE (1815), partial (also others)

Phaenopoda of MAYER (1824), partial (also “anilioids”)

Boidæ of GRAY (1825), approximate

Innocua of HAWORTH (1825), partial

Pythonoidea of FITZINGER (1826), approximate

Onychophori of RITGEN (1826), partial (also “anilioids”)

Macrostomata of MÜLLER (1831), partial (also others)

Peropodes of WIEGMANN (1832), approximate

Boas of SCHLEGEL (1837), partial (also acrochordids)

Saurophidia of FITZINGER (1843), partial (also scolecophidians and “anilioids”)

Azémiophides of DUMÉRIL & BIBRON (1844), partial (also “anilioids”)

Aglyphodontes of DUMÉRIL (1853), partial (also “anilioids” and others)

Asinea of COPE (1864), approximate

Boidiens of JAN (1865), approximate

Boaeformes of HEILPRIN (1907), approximate

Henophidia of HOFFSTETTER (1939), partial (also “anilioids” and others)

Booidea of GAUTHIER *et al.* (2012), approximate

Henophidia of WALLACH *et al.* (2014), approximate

### Diagnoses of Constrictores, Pythonoidea, and Booidea

A number of features have previously been listed as capable of differentiating “pythons” (Pythonidae sensu WALLACH *et al.*, 2014) and “boas” (Booidea sensu PYRON *et al.*, 2014). Whereas many of these features are serviceable in the sense of a dichotomous key, in many cases either the characters have not been included, or character polarity is ambiguous when evaluated in, broad-scale studies of snake phylogeny (e.g., GAUTHIER *et al.*, 2012). The most important features lie in the cranial anatomy, and some of them were already recognized in the 19<sup>th</sup> century. Pythonids (as well as *Loxocemus bicolor* Cope, 1861) possess premaxillary teeth (with the exception of the Australian genus *Aspidites* Peters, 1877), in contrast to boids, where the premaxilla is always toothless (FRAZZETTA, 1975; SZYNDLAR & RAGE, 2003). Furthermore, in pythonids, a foramen is present in the palatine for the infraorbital nerve

of maxillary division of the trigeminal, a feature that is absent in booids (SZYNDLAR & RAGE, 2003). Also, pythonids (as well as *Loxocemus bicolor*) possess a supraorbital bone, in contrast to booids (except *Calabaria reinhardtii* [Schlegel, 1851]), which lack it (ROMER, 1956; FRAZZETTA, 1975). On the basis of vertebral anatomy, boids share strong resemblance with pythonids; both have the massively built vertebrae, with a generally low ratio of their centrum length / neural arch width (<1.1), the high neural spines, and a thick zygosphene (IVANOV, 2000; SZYNDLAR & RAGE, 2003; GEORGALIS & SCHEYER, 2019). Nevertheless, Boidae (sensu PYRON *et al.*, 2014) usually have paracotylar foramina on their vertebrae, whereas pythonids and the majority of non-boid booids almost always lack them (KLUGE, 1993; SZYNDLAR & SCHLEICH, 1993; SZYNDLAR & RAGE, 2003; GEORGALIS, 2019; pers. observ.). However, this character can be variable and indeed its taxonomic utility has been questioned (RAGE, 2001). Furthermore, in pythonids, the shape of the haemal keel is defined by grooves or depressions beginning at the cotylar rim, but projecting below the centrum only in the posterior part of each vertebra (SCANLON & MACKNESS, 2002; SZYNDLAR & RAGE, 2003). Pythonids usually possess a higher number of vertebrae in comparison with booids (SCHAAL, 2004), but SCANFERLA & SMITH (2020) recently showed that some extinct booids had as many vertebrae as pythonids. Also, it can be stated that large pythonids possess thicker zygosphenes in comparison with similarly sized booids, but this is also subjected to variability (GLG, pers. obs.). Finally, pythonids are usually characterized by a relatively homogeneous intracolumnar vertebral morphology, in contrast to booids (SZYNDLAR & RAGE, 2003).

Modern diagnoses consistent with a current understanding of relationships have not been provided for Constrictores, Booidea or Pythonoidea. To determine morphological apomorphies diagnostic of those clades that are compatible with the reference phylogeny, we took the morphological data matrices of HSIANG *et al.* (2015, hereafter HEA), for osteology, and REEDER *et al.* (2015, hereafter REA), for squamation, and subjected them to maximum parsimony analysis in PAUP, using the phylogenetic tree of BURBRINK *et al.* (2020:Data File S7) as a backbone topological constraint. In both matrices, we took all anguimorph taxa as outgroups and deleted Gekkota, Dibamidae, Scinciformata, Laterata, Iguania, and Rhynchocephalia. Because the primary reference phylogeny is based on molecular data and the content of Constrictores in morphological analyses is different (lacking *Xenopeltis*, *Loxocemus*, and Bolyeriidae), as noted above, we refrain from providing diagnoses based on such topologies. For diagnoses of Constrictores, Pythonoidea and Booidea for a tree in which Bolyeriidae and *Xenophidion* fall outside Constrictores, see SCANFERLA & SMITH (2020:Document S1, section 2.3).

Unambiguous synapomorphies (i.e., those character state changes optimized under both acctrans and deltrans as synapomorphies of the clade in question) are as follows. Note that Bolyeriidae is not considered to belong either to Booidea or to Pythonoidea. In the reference phylogeny

(BURBRINK *et al.*, 2020) it is the immediate sister-group to Booidea, but if its position were to shift some of these diagnostic features might change.

**Constrictores.** Maxillary process of premaxilla tapers to a point distally (HEA 5/1); dorsum sellae enclosed in distinct fossa (HEA 404/2); Vidian canal caudal opening within basisphenoid (HEA 430/0); dentary mental foramen position displaced caudally (HEA 470/1). Additionally, we note that anteroposteriorly short vertebrae – that is, those with a low centrum length / neural arch width ratio (< 1.1) are associated with this clade. The ratio is higher in most outgroups and lower in all ingroup taxa (including Bolyeriidae, based on figs. 1–2 in HECHT & LADUKE, 1988) except Ungaliophiinae and *Xenopeltis* (e.g., SMITH, 2013). However, the vertebrae of *Xenophidion* have not been described (cf. WALLACH & GÜNTHER, 1998) and the taxon is not included in the reference phylogeny of BURBRINK *et al.* (2020). Moreover, *Xenopeltis* with elongate vertebrae (SMITH, 2013) is basal in Pythonoidea. Finally, the character is not unique to Constrictores, as it also occurs in Madtsoiidae and certain other snake taxa like *Acrochordus* (see HOFFSTETTER & GAYRARD, 1964; ZAHER *et al.*, 2019) and Tropidophiidae (see BOGERT, 1968a).

**Pythonoidea.** Medial frontal pillar suture to subolfactory process (HEA 54/2); ectopterygoid overlap of pterygoid long (HEA 362/1); posterior auditory foramen enclosed entirely in prootic (HEA 393/1); splenial anterior inferior alveolar foramen absent (HEA 491/1); premaxillary teeth absent on midline but present at lateral margins of element (HEA 547/1); elongate postmentals present (REA 650/1).

**Booidea.** Premaxilla internasal process narrowly clasped between nasals (HEA 15/2); frontal descending process abuts parietal (HEA 56/2); suture between frontal and parietal in medial wall of orbit vertical or only slightly anteriorly inclined (HEA 69/1); frontal suboptic shelf below optic foramen deep (HEA 72/1); quadrate suprastapedial process absent (HEA 219/0); lateral edge of stapedial footplate nearly in same cross-sectional plane as medial edge (HEA 238/1); posterior base of lateral flange of septomaxilla distinctly cranial to vomeronasal organ (HEA 252/1); palatine maxillary process at posterior end of palatine (HEA 297/1); ectopterygoid maxillary process tapering or parallel-sided (HEA 347/0); ectopterygoid abuts pterygoid laterally (HEA 358/2); cranial rim of crista circumfenestralis caudal extent relative to medial margin of stapedial footplate: former roughly on same level as latter in dorsal view at level of shaft (HEA 385/1); maxillary branch of trigeminal nerve passes dorsally between palatine and prefrontal (HEA 401/1); posterior opening of right Vidian canal large (HEA 422/1); angular process of dentary terminates well posterior to splenio-angular joint (HEA 476/2); coronoid eminence composed of both surangular and coronoid (HEA 501/0); retroarticular process (*in situ*) extends posteriorly (HEA 529/0); dentary teeth conspicuously enlarged anteriorly (HEA 544/1).

## Acknowledgments

We thank Pr. Patrick Smith, Thomas Lehmann, Kevin de Queiroz, and Van Wallach for discussion and Nico Cellinese for help with registration. GLG acknowledges support from a postdoctoral grant of the Università degli Studi di Torino and the National Scholarship Program of the Slovak Republic (SAIA). GLG also acknowledges study of comparative material through the grants SYNTHESYS ES-TAF-5910 (MNCN), SYNTHESYS AT-TAF-5911 (NHMW), SYNTHESYS HU-TAF-6145 (HNHM), and SYNTHESYS GB-TAF-6591 (NHMUK), and the respective curators (Marta Calvo-Revuelta, Silke Schweiger, Judit Vörös, and Sandra Chapman) are highly thanked here. For permission to use their photographs of extant Constrictores in Fig. 1, we are grateful to Ilias Strachinis and Alberto Sanchez Vialas. The quality of the manuscript was enhanced by useful comments made by the Editor Uwe Fritz and two anonymous reviewers. We also thank anonymous reviewers of a previous version of this manuscript, who pointed out overlooked literature and useful comments that improved the paper.

## References

- BASZIO, S. (2004). *Messelophis variatus* n.gen. n.sp., from the Eocene of Messel: a tropidopheine snake with affinities to Erycinae (Boidae). *Courier Forschungsinstitut Senckenberg*, **252**, 47–66.
- BEDRIAGA, J. VON (1882). Die Amphibien und Reptilien Griechenlands. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **56**, 1–195.
- BOBACK, S. M., McCANN, K. J., WOOD, K. A., MCNEAL, P. M., BLANKENSHIP, E. L. & ZWEMER, C. F. (2015). Snake constriction rapidly induces circulatory arrest in rats. *The Journal of Experimental Biology*, **218**, 2279–2288.
- BODDAERT, P. (1783). Specimen novae methodi distinguendi Serpentia. *Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum*, **7**, 12–27.
- BOETTGER, O. (1884). III. Systematik, Faunistik, Biologie. 3. Reptilia. *Zoologischer Jahresbericht*, **1884**, 214–261.
- BOGERT, C. M. (1968a). A new genus and species of dwarf boa from Southern Mexico. *American Museum Novitates*, **2354**, 1–38.
- BOGERT, C. M. (1968b). The variations and affinities of the dwarf boas of the genus *Ungaliophis*. *American Museum Novitates*, **2340**, 1–26.
- BOIÉ, F. (1826). Generalübersicht der Familien und Gattungen der Ophidier. *Isis von Oken, Leipzig*, **19**, 981–982.
- BOIÉ, F. (1827). Bemerkungen über Merrem's Versuch eines Systems der Amphibien, 1-ste Lieferung: Ophidier. *Isis von Oken, Leipzig*, **20**, 508–566.
- BONAPARTE, C. L. (1831). Saggio d'una distribuzione metodica degli animali vertebrati a sangue freddo. *Giornale Arcadico di Scienze, Lettere, ed Arti, Roma*, **52**, 129–209.
- BONNATERRE, J. P. (1790). *Ophiologie*. In: *Tableau encyclopédique et méthodique des trois règnes de la nature*. Paris, Panconoe, 76 pp.
- BOULENGER, G. A. (1890). *The Fauna of British India, including Ceylon and Burma. Reptilia and Batrachia*. London, Taylor and Francis, 541 pp.
- BOULENGER, G. A. (1893). *Catalogue of the Snakes in the British Museum (Natural History). Volume 1, Containing the Families Typhlopidae, Glauconidae, Boidae, Ilysiidae, Uropeltidae, Xenopeltidae, and Colubridae Aglyphae, part*. London, Trustees of the British Museum (Natural History), 448 pp.

- BOULENGER, G. A. (1913). *The Snakes of Europe*. London, Methuen & Company Limited, 269 pp.
- BREHM, A. E. (1878). *Merveilles de la nature: l'homme et les animaux, description populaire des races humaines et des règne animal*. Translation in French by H.E. Sauvage. Paris, J.-B. Bailliére et Fils, 726 pp.
- BURBRINK, F. T. & CROTHER, B. I. (2011). Evolution and taxonomy of snakes, pp. 19–53 in ALDRIDGE, R. A. & SEVER, D. M. (eds) *Reproductive Biology and Phylogeny of Snakes*. Boca Raton, CRC Press.
- BURBRINK, F. T., GRAZZIOTIN, F. G., PYRON, R. A., CUNDALL, D., DONNELLAN, S., IRISH, F., KEOGH, J. S., KRAUS, F., MURPHY, R. W., NOONAN, B., RAXWORTHY, C. J., RUANE, S., LEMMON, A. R., LEMMON, E. M. & ZAHER, H. (2020). Interrogating genomic-scale data for Squamata (lizards, snakes, and amphisbaenians) shows no support for key traditional morphological relationships. *Systematic Biology*, **69**, 502–520.
- BURMEISTER, H. (1837). *Handbuch der Naturgeschichte. Zum Gebrauch bei Vorlesungen*. Berlin, T.C.F. Enslin, 858 pp.
- CANTINO, P. D. & DE QUEIROZ, K. (2014). *International Code of Phylogenetic Nomenclature, version 5*. Available at: <https://www.ohio.edu/phylocode/>
- CANTOR, T. E. (1847). *Catalogue of Reptiles inhabiting the Malayan Peninsula and Islands, Collected or Observed by Theodore Cantor, Esq., M.D. Bengal Medical Service*. Calcutta, J. Thomas, 157 pp.
- CARUS, J. V. (1868). *Handbuch der Zoologie. Wirbelthiere*. Leipzig, W. Engelmann, 432 pp.
- COPE, E. D. (1861). "Some remarks defining ... species of Reptilia Squamata..." *Proceedings of the Academy of Natural Sciences of Philadelphia*, **13**, 73–77.
- COPE, E. D. (1862). Notes upon some reptiles of the Old World. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **14**, 337–344.
- COPE, E. D. (1864). On the characters of the higher Groups of Reptilia Squamata – and especially of Diploglossa. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **15**, 224–231.
- COPE, E. D. (1883). *The Vertebrata of the Tertiary Formations of the West (Report of the United States Geological Survey of the Territories, Vol. III)*. Washington D.C., Government Printing Office, 1009 pp.
- COPE, E. D. (1886). An analytical table of the genera of snakes. *Proceedings of the American Philosophical Society held at Philadelphia for promoting useful knowledge*, **23**, 479–499.
- COPE, E. D. (1893). Prodromus of a new System of the non-venomous snakes. *The American Naturalist*, **27**, 477–483.
- COPE, E. D. (1894). The classification of snakes. *The American Naturalist*, **28**, 831–844.
- COPE, E. D. (1895). The classification of the Ophidia. *Transactions of the American Philosophical Society*, **18**, 186–219.
- COPE, E. D. (1898). *Syllabus of lectures on the vertebrata*. Philadelphia, University of Pennsylvania, 135 pp.
- CUNDALL, D. & IRISH, F. J. (1986). Aspects of locomotor and feeding behavior in the Round Island boa, *Casarea dussumieri*. *Dodo*, **23**, 108–111.
- CUNDALL, D., WALLACH, V. & ROSSMAN D. A. (1993). The systematic relationships of the snake genus *Anomochilus*. *Zoological Journal of the Linnean Society*, **109**, 275–299.
- CUVIER, G. (1817). *Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Tome II, contenant les reptiles, les poissons, les mollusques et les annélides*. Paris, Deterville, 532 pp.
- CUVIER, G. (1829). *Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Tome II. Nouvelle édition, revue et augmentée*. Paris, Deterville, 406 pp.
- DAUDIN, F. M. (1803a). Erpétologie. Caractères des vingt-trois genres qui composent l'ordre des Ophidiens. *Magasin Encyclopédique, ou Journal des Sciences, des Lettres et des Arts, Paris*, **5**, 433–438.
- DAUDIN, F. M. (1803b). *Histoire naturelle, générale et particulière des reptiles; ouvrage faisant suite aux Oeuvres de Leclerc de Buffon, et partie du cours complet d'histoire naturelle rédigé par C.S. Sonnini, membre de plusieurs sociétés savantes. Tome cinquième*. Paris, F. Dufart, 365 pp.
- DAUDIN, F. M. (1803c). *Histoire naturelle, générale et particulière des reptiles; ouvrage faisant suite aux Oeuvres de Leclerc de Buffon, et partie du cours complet d'histoire naturelle rédigé par C.S. Sonnini, membre de plusieurs sociétés savantes. Tome sixième*. Paris, F. Dufart, 447 pp.
- DAUDIN, F. M. (1803d). *Histoire naturelle, générale et particulière des reptiles: ouvrage faisant suite à l'histoire naturelle générale et particulière, composée par Leclerc de Buffon, et rédigée par C.S. Sonnini. Tome septième*. Paris, F. Dufart, 436 pp.
- DAY, D. (1989). *The doomsday book of animals. A natural history of vanished species*. New York, Viking Press, 288 pp.
- DIESING, K. M. (1851). *Systema helminthum. Vindobonae (= Viena)*, W. Braumüller, 588 pp.
- DOWLING, H. G., HASS, C. A., HEDGES, S. B. & HIGHTON, R. (1996). Snake relationships revealed by slow-evolving proteins: a preliminary survey. *Journal of Zoology, London*, **240**, 1–28.
- DUMÉRIL, A. (1805). *Analytische Zoologie. Aus der Französischen, mit Zusätzen, von L. F. Froriep*. Weimar, Verlag des Landes-Industrie-Comptoirs, 544 pp.
- DUMÉRIL, A. (1853). Prodrome de la classification des reptiles ophiidiens. *Mémoires de l'Académie scientifique de Paris*, **23**, 399–535.
- DUMÉRIL, A. (1859). Reptiles et Poissons de l'Afrique occidentale. Etude precedee de considerations générales sur leur distribution géographique. *Archives du Muséum National d'Histoire Naturelle, Paris*, **10**, 137–240.
- DUMÉRIL, A. M. C. & BIBRON, G. (1844). *Erpétologie générale ou histoire naturelle complète des reptiles. Tome sixième, comprenant l'histoire générale des ophiidiens, la description des genres et des espèces de serpents non venimeux, savoir, la totalité des vermiformes ou des scolopophides, et partie des circuiriformes ou azémiophides; en tout vingt-cinq genres et soixante-cinq espèces*. Paris, Librairie Encyclopédique de Roret, 609 pp.
- DUMÉRIL, A. M. C., BIBRON, G. & DUMÉRIL, A. (1854a). *Erpétologie générale ou histoire naturelle complète des reptiles. Tome septième – première partie. Compenant l'histoire des serpentes non venimeux*. Paris, Librairie Encyclopédique de Roret, 780 pp.
- DUMÉRIL, A. M. C., BIBRON, G. & DUMÉRIL, A. (1854b). *Erpétologie générale ou histoire naturelle complète des reptiles. Tome neuvième. Contenant Contenant l'histoire des batraciens, urodèles. Plus, un répertoire ou catalogue méthodique de tous les reptiles décrits dans les neuf volumes; avec des tables générales pour le texte et pour les figures*. Paris, Librairie Encyclopédique de Roret, 440 pp.
- EICHWALD, K. E. I. (1831). *Zoologia specialis quam expositis animalibus tum vivis, tum fossilibus potissimum Rossiae in universum, et Poloniae in specie, in usum lectionum publicarum in Universitate Caesarea Vilnensi habendarum. Pars posterior [vol.3], specialem expositionem spondylozoorum continens*. Vilnius, Josephi Zawadzki, 369 pp.
- FAYRER, J. (1872). *The Thanatophidia of India, being a Description of the Venomous Snakes of the Indian Peninsula, with an Account of the Influence of their Poison on Life; and a Series of Experiments*. London, J. & A. Churchill, 131 pp.
- FAYRER, J. (1877). Venomous animals. *Edinburgh Medical Journal*, **23**, 97–117.
- FEDOROVICH GORIANINOW, P. (1834). *Primae lineae systematis naturae: nexui naturali omnium evolutionique progressivae per nixus reascendentes superstructi*. Petropoli (Saint Petersburg), Karolis Krajanis, 142 pp.
- FIGUEROA, A., MCKELVY, A. D., GRISMER, L. L., BELL, C. D. & LAILVAUX, S. P. (2016). A species-level phylogeny of extant snakes

- with description of a new colubrid subfamily and genus. *PLoS One*, **11**, e0161070.
- FILHOL, H. (1877). Recherches sur les Phosphorites du Quercy. Étude des fossiles qu'on y rencontre et spécialement des mammifères. Pt. II. *Annales des Sciences géologiques*, **8**, 1–340.
- FILIPPI, F. DE (1840). *Catalogo ragionato e descrittivo della raccolta de' serpenti del Museo dell'I.R. Università di Pavia*. Milan, Biblioteca Italiana, 65 pp.
- FISCHER VON WALDHEIM, G. (1813). *Zoognosia tabulis synopticas illustrata, in usum praelectionum Academiae Imperialis Medico-Chirurgicae Mosquensis edita. Editio tertia, classium, ordinum, generum illustratione perpetua aucta. Volumen primum, tabulis synopticas generales et comparativas, nec non characteram quorundam explicationem iconographicam continens. Editio tertia, classium, ordinum, generum illustratione perpetua aucta. Volumen primum, tabulis synopticas generales et comparativas, nec non characterum quorundam explicationem iconographicam continens. Parte tertia (Reptiles, poisons)*. Moscow, Nicolai Sergeidis Vsevolozsky, pp. 57–117.
- FITZINGER, L. J. F. J. (1826). *Neue Classification der Reptilien nach ihren Natürlichen Verwandtschaften. Nebst einer Verwandschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des k. k. zoologischen Museums zu Wien*. Wien, J. G. Huebner, 66 pp.
- FITZINGER, L. J. F. J. (1843). *Systema reptilium. Fasciculus primus. Amblyglossae*. Vindobonae (= Vienna), Braumüller et Seidel Bibliopolas, 106 pp.
- FITZINGER, L. J. F. J. (1861). Die Ausbeute der österreichischen Naturforscher an Säugetieren und Reptilien während der Weltumsegelung Sr. Majestät Fregatte Novara. Novara. *Sitzungsberichte der Akademie der Wissenschaften in Wien, Mathematische-Naturwissenschaftliche Klasse*, **42**, 383–416.
- FITZINGER, L. J. F. J. (1867). *Bilder-Atlas zur wissenschaftlich-populären Naturgeschichte der Wirbelthiere*. Vienna, Verlag der Kaiserlich-Königlichen Hof- und Staatsdruckerei, 102 pp.
- FRASER, A. G. L. (1937). The snakes of Deolali. With notes on their comparative osteology and peculiarities of dentition. Part III. *Journal of the Bombay Natural History Society*, **39**, 464–501.
- FRAZZETTA, T. H. (1959). Studies on the morphology and function of the skull in the Boidae (Serpentes). Part 1. Cranial differences between *Python sebae* and *Epicrates cenchria*. *Bulletin of the Museum of Comparative Zoology, Harvard University*, **119**, 453–472.
- FRAZZETTA, T. H. (1966). Studies on the morphology and function of the skull in the Boidae (Serpentes). Part 2. Morphology and function of the jaw apparatus in *Python sebae* and *Python molurus*. *Journal of Morphology*, **118**, 217–296.
- FRAZZETTA, T. H. (1975). Pattern and instability in the evolving premaxilla of boine snakes. *American Zoologist*, **15**, 469–481.
- GADOW, H. (1909). *Amphibia and Reptilia*. London, MacMillan, 668 pp.
- GASC, J.-P. (1974). L'interprétation fonctionnelle de l'appareil musculo-squelettique de l'axe vertébral chez les serpents (Reptilia). *Mémoires du Muséum National d'Histoire Naturelle, Paris, Série A*, **83**, 1–182.
- GAUTHIER, J. A., KEARNEY, M., MAISANO, J. A., RIEPPEL, O. & BEHKE, A. D. B. (2012). Assembling the squamate tree of life: perspectives from the phenotype and the fossil record. *Bulletin of the Peabody Museum of Natural History*, **53**, 3–308.
- GEORGALIS, G. L. (2019). Poor but classic: The squamate fauna from the late Miocene of Pikermi, near Athens, Greece. *Comptes Rendus Palevol*, **18**, 801–815.
- GEORGALIS, G. L., DEL FAVERO, L. & DELFINO, M. (2020). Italy's largest snake - redescription of *Palaeophis oweni* from the Eocene of Monte Duello, near Verona. *Acta Palaeontologica Polonica*, **65**,
- GEORGALIS, G. L. & SCHAYER, T. M. (2019). A new species of *Palaeopython* (Serpentes) and other extinct squamates from the Eocene of Dielsdorf (Zurich, Switzerland). *Swiss Journal of Geosciences*, **112**, 383–417.
- GILMORE, C. W. (1938). Fossil snakes of North America. *Geological Society of North America, Special Papers*, **9**, 1–96.
- GIRARD, H. (1895). *Aide-mémoire de Zoologie*. Paris, J.-B. Bailière, 300 pp.
- GMELIN, J. F. (1789). *Caroli a Linne...Systema naturae per regna tria naturalia, secundum classes, ordines, genera, species, cum characteribus differentiis, synonymis, locis. Tomus I, Editio decima tertia, aucta, reformata. Pars III. Amphibia et Pisces*. Leipzig, Georg Emanuel Beer, pp. 1038–1516.
- GORSKI, C. (1852). *Ueber das Becken der Saurier: eine vergleichend-anatomische Abhandlung*. Dorpat, H. Laakmann, 48 pp.
- GOWER, D. J., VIDAL, N., SPINKS, J. N. & McCARTHY, C. J. (2005). First inclusion of Anomochilidae in molecular phylogenetics of the major lineages of snakes. *Journal of Zoological Systematics and Evolutionary Research*, **43**, 315–320.
- GRAY, J. E. (1825). A synopsis of the genera of Reptiles and Amphiibia, with a description of some new species. *Annals of Philosophy, Series 2*, **10**, 193–217.
- GRAY, J. E. (1842). Synopsis of the species of prehensile-tailed snakes, or family Boidæ. *Zoological Miscellany*, **2**, 41–46.
- GRAY, J. E. (1849). *Catalogue of the Specimens of Snakes in the Collection of the British Museum*. London, British Museum (Natural History), 125 pp.
- GRAY, J. E. (1858). Description of a new genus of Boidæ from Old Calabar, and a list of W. African reptiles. *Proceedings of the Zoological Society of London*, **26**, 154–167.
- GROOMBRIDGE, B. C. (1979). Variations in morphology of the superficial palate of henophidian snakes and some possible systematic implications. *Journal of Natural History*, **13**, 661–680.
- GÜNTHER, A. C. L. G. (1864). *The Reptiles of British India*. London, Ray Society, 452 pp.
- GÜNTHER, A. C. L. G. (1877). On a collection of reptiles and fishes from Duke of York Island, New Ireland, and New Britain. *Proceedings of the Zoological Society of London*, **1877**, 127–132.
- GÜNTHER, R. & MANTHEY, U. (1995). *Xenophidion*: a new genus with two new species of snakes from Malaysia (Serpentes, Colubridae). *Amphibia-Reptilia*, **16**, 229–240.
- HAAS, G. (1952). The head muscles of the genus *Causus* (Ophidia, Solenoglypha) and some remarks on the origin of the Solenoglypha. *Proceedings of the Zoological Society of London*, **122**, 573–592.
- HARDING, J. H. & HOLMAN, J. A. (1981). The paleohistory of snakes in North America - A brief review. *Bulletin of the Chicago Herpetological Society*, **17**, 33–45.
- HARRINGTON, S. M. & REEDER, T. W. (2017). Phylogenetic inference and divergence dating of snakes using molecules, morphology and fossils: new insights into convergent evolution of feeding morphology and limb reduction. *Biological Journal of the Linnean Society*, **121**, 379–394.
- HAWORTH, A. H. (1825). A binary arrangement of the Class Amphiibia. *Philosophical Magazine and Journal*, **65**, 372–373.
- HEAD, J. J. (2015). Fossil calibration dates for molecular phylogenetic analysis of snakes 1: Serpentes, Alethinophidia, Boidae, Pythonidae. *Palaeontologia Electronica*, **18**, 1.6FC.
- HEAD, J. J., BLOCH, J. I., HASTINGS, A. K., BOURQUE, J. R., CADENA, E. A., HERRERA, F. A., POLLY, P. D. & JARAMILLO, C. A. (2009). Giant boid snake from the Paleocene neotropics reveals hotter past equatorial temperatures. *Nature*, **457**, 715–718.
- HECHT, M. K. (1959). Amphibians and Reptiles in: McGREW, P. O., BERMAN, J. E., HECHT, M. K., HUMMEL, J. M., SIMPSON, G. G. & WOOD, A. E. (eds) The geology and paleontology of the Elk Mountain and Tabernacle Butte area, Wyoming. *Bulletin of American Museum of Natural History*, **117**, 130–146.
- HECHT, M. K. & LADUKE, T. C. (1988). Bolyerine vertebral variation: A problem for paleoherpetology. *Acta Zoologica Cracoviensis*, **31**, 605–614.
- HEILPRIN, A. (1907). The geographical and geological distribution of animals. Third Edition. *The International scientific series*, **57**, 1–435.

- HOFFMANN, C. K. (1890). *Dr. H.G. Bronn's Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild. Sechster Band. III. Abtheilung. Reptilien. III. Schlangen und Entwicklungsgeschichte der Reptilien.* Leipzig, Heidelberg, C. F. Winter, pp. 1401–2089.
- HOFFSTETTER, R. (1939). Contribution à l'étude des Elapidæ actuels et fossiles et de l'ostéologie des Ophidiens. *Archives du Muséum d'Histoire Naturelle de Lyon*, **15**, 1–78.
- HOFFSTETTER, R. (1946). Remarques sur la classification des Ophidiens et particulièrement des Boides des Mascareignes (Boyerinae subfam. nov.). *Bulletin du Muséum National d'Histoire Naturelle, Série 2*, **18**, 132–135.
- HOFFSTETTER, R. (1955). Squamates de type moderne, pp. 606–662 in: PIVETEAU, J. (ed.) *Traité de Paléontologie, Vol. 5.* Paris, Masson.
- HOFFSTETTER, R. & GAYRARD, Y. (1965). Observations sur l'ostéologie et la classification des Acrochordidae (Serpentes). *Bulletin du Muséum National d'Histoire Naturelle, Paris, Série 2*, **36**, 677–696.
- HOFFSTETTER, R. & RAGE, J.-C. (1972). Les Erycinæ fossiles de France (Serpentes, Boidæ). Compréhension et histoire de la sous-famille. *Annales de Paléontologie*, **58**, 81–124.
- HOLMAN, J. A. (2000). *Fossil snakes of North America: Origin, Evolution, Distribution, Paleoecology.* Bloomington & Indianapolis, Indiana University Press, 357 pp.
- HORNSTEDT, C. F. (1787). Beskrifning po en ny orm från Java. *Kongliga Svenska Vetenskaps-Akademiens Handlingar, Stockholm*, **8**, 306–308.
- HSIANG, A. Y., FIELD, D. J., WEBSTER, T. H., BEHLKE, A. D. B., DAVIS, M. B., RACICOT, R. A. & GAUTHIER, J. A. (2015). The origin of snakes: revealing the ecology, behavior, and evolutionary history of early snakes using genomics, phenomics, and the fossil record. *BMC Evolutionary Biology*, **15**, 87.
- [ICZN] INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE (1999). *International Code of Zoological Nomenclature. 4<sup>th</sup> ed.* London, International Trust for Zoological Nomenclature, 306 pp.
- IHERING, R. VON (1911). As Cobras do Brasil. *Revista do Museu Paulista*, **8**, 273–378.
- IVANOV, M. (2000). Snakes of the lower/middle Miocene transition at Vieux Collonges (Rhône; France), with comments on the colonisation of western Europe by colubroids. *Geodiversitas*, **22**, 559–588.
- JAN, G. (1862). Ueber die Familien der Eryciden und Tortriciden. *Archiv für Naturgeschichte*, **28**, 238–252.
- JAN, G. (1863). *Elenco sistematico degli ofidi descritti e disegnati per l'Iconografia générale.* Milan, A. Lombardi, 143 pp.
- JAN, G. (1865). *Iconographie générale des ophidiens. Deuxième famille, les Uropeltiens; troisième famille, les Tortriciens; quatrième famille, les Boidiens.* Paris, J. B. Baillière et fils, pp. 43–100.
- JAN, G. & SORDELLI, F. (1860–1866). *Iconographie générale des ophidiens. Tome premier (livrais 1 à 17), contenant cent deux planches.* Milan, Georges Jan & Ferdinand Sordelli.
- KIRITZESCU, C. (1902). Sur la présence d'*Eryx jaculus* en Roumanie. *Buletinul Societății de Științe din București, România*, **11**, 620–626.
- KLUGE, A. G. (1993). Calabaria and the phylogeny of erycine snakes. *Zoological Journal of the Linnean Society*, **107**, 293–351.
- KUHN, O. (1946). Das System der fossilen und rezenten Amphibien und Reptilien. *Bericht der Naturforschenden Gesellschaft in Bamberg*, **29**, 49–67.
- KUHN, O. (1963). *Fossilium Catalogus, Volume I: Animalia, part 103, Serpentes (Supplementum I).* The Hague, W. Junk, 45 pp.
- LACÉPÈDE, B. G. E. (1789). Histoire naturelle des quadrupèdes ovipares et des serpens. Tome second (Histoire naturelle des serpens). Hôtel de Thou, Paris, (20) + 144 pp. (Table méthodique) + 527 pp.
- LACÉPÈDE, B. G. E. (1801). Sur un nouveau genre de serpent. *Bulletin de la Société Philomathique de Paris, Série 2*, **46**, 169.
- LATREILLE, P. A. (1804). *Familles naturelles du règne animal, exposées succinctement et dans un ordre analytique, avec l'indication de leurs genres.* Paris, J.B. Bailliére, 570 pp.
- LAWSON, R., SLOWINSKI, J. B. & BURBRINK, F. T. (2004). A molecular approach to discerning the phylogenetic placement of the enigmatic snake *Xenophidion schaeferi* among the Alethinophidia. *Journal of Zoology, London*, **263**, 1–10.
- LEBLANC, E. (1920). *Recherches sur les plexus choroides des reptiles.* Alger, Jules Carbonel, 123 pp.
- LEE, M. S. Y., HUGALL, A. F., LAWSON, R. & SCANLON, J. D. (2007). Phylogeny of snakes (Serpentes): combining morphological and molecular data in likelihood, Bayesian and parsimony analyses. *Systematics and Biodiversity*, **5**, 371–389.
- LEE, M. & SCANLON, J. D. (2002). Snake phylogeny based on osteology, soft anatomy and behaviour. *Biological Reviews*, **77**, 333–402.
- LEUCKART, F. S. (1841). *Zoologische Bruchstücke. Vol. 2.* Stuttgart, F. L. Rieger & Comp., 130 pp.
- LI, J. N., LIANG, D., WANG, Y.-Y., GUO, P., HUANG, S. & ZHANG, P. (2020). A large-scale systematic framework of Chinese snakes based on a unified multilocus marker system. *Molecular Phylogenetics and Evolution*, **148**, 106807.
- LICHENSTEIN, M. H. C. & VON MARTENS, E. (1856). *Nomenclator reptilium et amphibiorum Musei Zoologici Berolinensis.* Berlin, Königliche Akademie der Wissenschaften, 48 pp.
- LINK, H. F. (1807). *Beschreibung der Naturalien-Sammlung der Universität zu Rostock. Zweite Abtheilung.* Rostock, Adlers Erben, pp. 51–100.
- LINNAEUS, C. (1758). *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis.* Stockholm, Laurentius Salvius, 824 pp.
- LYDEKKER, R. (1888). *Catalogue of Fossil Reptiles and Amphibians in the British Museum (Natural History). Part I. Containing the Orders Ornithosauria, Crocodilia, Dinosauria, Squamata, Rhynchocephalia, and Proterosauria.* London, British Museum (Natural History), 309 pp.
- MAYER, DE B. (1824). Über die hintere Extremität der Ophidien. *Nova acta physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosum*, **12**, 821–842.
- MCDOWELL, S. B. (1987). Systematics, pp. 3–50 in: SEIGEL, R. A., COLLINS, J. T. & NOVAK, S. S. (eds.) *Snakes: Ecology and Evolutionary Biology.* New York, Macmillan.
- MEYER, A. B. (1874). Übersicht der von mir auf Neu-Guinea und den Inseln Jobi, Mysore und Mafoor im Jahre 1873 gesammelten Amphibien. *Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin*, **1874**, 128–140.
- MIRALLES, A., MARIN, J., MARKUS, D., HERREL, A., HEDGES, B. S. & VIDAL, N. (2018). Molecular evidence for the paraphyly of Scolecophidia and its evolutionary implications. *Journal of Evolutionary Biology*, **31**, 1782–1793.
- MÜLLER, J. (1831). Beiträge zur Anatomie und Naturgeschichte der Amphibien. *Zeitschrift für Physiologie*, **4**, 190–275.
- MÜLLER, F. (1878). Katalog der im Museum und Universitätskabinett zu Basel aufgestellten Amphibien und Reptilien nebst Anmerkungen. *Verhandlungen der Naturforschenden Gesellschaft in Basel*, **6**, 557–709.
- MÜLLER, F. (1880). Erster Nachtrag zum Katalog der herpetologischen Sammlung des Basler Museums. *Verhandlungen der Naturforschenden Gesellschaft in Basel*, **7**, 120–165.
- MURPHY, J. C. & HENDERSON, R. W. (1997). *Tales of Giant Snakes: A Historical Natural History of Anacondas and Pythons.* Malabar, Florida, Krieger Publishing, 221 pp.
- NICHOLSON, H.A. & LYDEKKER, R. (1889). *A Manual of Palaeontology, for the Use of Students; With a General Introduction on the Principles of Palaeontology. Volume II. Part III – Palaeozoology.* Edinburgh, W. Blackwood & Sons, 1624 pp.
- NOGUCHI, H. (1909). *Snake Venoms, an Investigation of Venomous Snakes with Special Reference to the Phenomena of their Venoms.* Washington, D.C., Carnegie Institution, 315 pp.

- NOONAN, B. P. & CHIPPINDE, P. T. (2006). Dispersal and vicariance: The complex evolutionary history of boid snakes. *Molecular Phylogenetics and Evolution*, **40**, 347–358.
- NOPCSA, F. (1923). *Eidolosaurus und Pachyophis*. Zwei neue Neocom-Reptilien. *Palaeontographica*, **65**, 99–154.
- OGUIURA, N., FERRAREZZI, H., & BATISTIC, R. F. (2010). Cytogenetics and molecular data in snakes: a phylogenetic approach. *Cytogenetics and Genome Research*, **127**, 128–142.
- OPPEL, M. (1811a). Suite du 1er. mémoire sur la classification des reptiles. Ord. II. Squammata mihi. Sect. II. Ophidii. Ord. III. Ophidii, Brongniart. *Annales du Muséum d'Histoire Naturelle, Paris*, **16**, 376–393.
- OPPEL, M. (1811b). *Die Ordnungen, Familien und Gattungen der Reptilien als Prodrom einer Naturgeschichte derselben*. Munich, Joseph Lindauer, 87 pp.
- PALACKÝ, J. (1884). Die Verbreitung der fossilen Schlangen in Europa. *Sitzungsberichte der Königlichen Böhmisches Gesellschaft der Wissenschaften in Prag*, **1884**, 165–166.
- PETERMANN, H. & GAUTHIER, J. A. (2018). Fingerprinting snakes: paleontological and paleoecological implications of zygantral growth rings in Serpentes. *PeerJ*, **6**, e4819.
- PETERS, W. C. H. (1860). Eine neue Gattung von Riesenschlangen, welche von einem gebornen Preußen, Hrn. Carl Reiss, in Guayaquil nebst mehreren anderen wertvollen Naturalien dem zoologischen Museum zugesandt worden ist. *Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin*, **1860**, 200–202.
- PETERS, W. C. H. (1877). Herpetologische Notizen. II. Bemerkungen über neue oder weniger bekannte Amphibien. *Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin*, **1877**, 415–423.
- PETERS, W. C. H. (1882). *Naturwissenschaftliche Reise nach Mosambique auf Befehl seiner Majestät des Königs Friedrich Wilhelm IV. In den Jahren 1842 bis 1848 ausgeführt von Wilhelm C. H. Peters. Zoologie. III. Amphibien*. Berlin, 191 pp.
- PYRON, R. A., BURBRINK, F. T. & WIENS, J. J. (2013). A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evolutionary Biology*, **13**, 93.
- PYRON, R. A., REYNOLDS, G. A. & BURBRINK, F. T. (2014). A taxonomic revision of boas (Serpentes: Boidae). *Zootaxa*, **3846**, 249–260.
- RAFINESQUE-SCHMALTZ, C. S. (1815). *Analyse de la nature ou tableau de l'univers et des corps organisés*. Palermo, Constantine Samuel Rafinesque-Schmaltz, 224 pp.
- RAGE, J.-C. (1977). An erycine snake (Boidae) of the genus *Calamagras* from the French lower Eocene, with comments on the phylogeny of the Eryciniae. *Herpetologica*, **33**, 459–463.
- RAGE, J.-C. (1984). *Encyclopedia of Paleoherpetology. Part 1, Serpentes*. Stuttgart & New York, Gustav Fischer, 80 pp.
- RAGE, J.-C. (2001). Fossil snakes from the Paleocene of São José de Itaboráí, Brazil. Part II. Boidae. *Palaeovertebrata*, **30**, 111–150.
- RAGE, J.-C., BAPAI, S., THEWISSEN, J. G. M. & TIWARI, B. N. (2003). Early Eocene snakes from Kutch, Western India, with a review of the Palaeophiidae. *Geodiversitas*, **25**, 695–716.
- RAMÓN DE LA SAGRA, D. (1838–1843). *Historia física, política y natural de la isla de Cuba. Tomo II. Segunda parte – Reptiles y peces. Reptiles [J.T. Cocteau and G. Bibron]*. Paris, Arthur Bertrand, 143 pp.
- REYNOLDS, R. G. & HENDERSON, R. W. (2018). Boas of the world (Superfamily Booidae): a checklist with systematic, taxonomic, and conservation assessments. *Bulletin of the Museum of Comparative Zoology*, **162**, 1–58.
- REYNOLDS, R. G., NIEMILLER, M. L. & REVELL, L. J. (2014). Toward a Tree-of-Life for the boas and pythons: Multilocus species-level phylogeny with unprecedented taxon sampling. *Molecular Phylogenetics and Evolution*, **71**, 201–213.
- RIEPPEL, O. (1977). Studies on the skull of the Henophidia (Reptilia: Serpentes). *Journal of Zoology, London*, **181**, 145–173.
- RIEPPEL, O. (1988). A review of the origin of snakes. *Evolutionary Biology*, **22**, 37–130.
- RITGEN, F.A.M.F. VON (1826). Versuch einer natürlichen Eintheilung der Amphibien. *Nova Acta Academiae Caesareae Leopoldino-Carolinae*, **14**, 246–284.
- ROCHEBRUNE, A. T. DE. (1880). Revision des ophidiens fossiles du Muséum d'Histoire Naturelle. *Nouvelles Archives du Muséum d'Histoire Naturelle, 2ème Série*, **3**, 271–296.
- ROCHEBRUNE, A. T. DE (1884). Faune ophiologique des Phosphorites du Quercy. *Mémoires de la Société des Sciences naturelles de Saône-et-Loire*, **5**, 149–164.
- ROMER, A. S. (1956). *Osteology of the Reptiles*. Chicago, University Chicago Press, 772 pp.
- ROUX-ESTÈVE, R. (1965). Les Serpents de la région de La Maboké - Boukoko. *Cahiers de la Maboké*, **3**, 51–92.
- SCANFERLA, C. A. & SMITH, K. T. (2020). Exquisitely preserved fossil snakes of Messel: insight into the evolution, biogeography, habitat preferences and sensory ecology of early boas. *Diversity*, **12**, 100.
- SCANFERLA, C. A., SMITH, K. T. & SCHAL, S. F. K. (2016). Revision of the cranial anatomy and phylogenetic relationships of the Eocene minute boas *Messelophis variatus* and *Messelophis ermannorum* (Serpentes, Booidea). *Zoological Journal of the Linnean Society*, **176**, 182–206.
- SCANLON, J. D. & LEE, M. S. Y. (2011). Chapter 3: The major clades of living snakes: morphological evolution, molecular phylogeny, and divergence dates, pp. 55–95 in: SEVER, D. M. & ALDRIDGE, R. D. (eds.) *Reproductive Biology and Phylogeny of Snakes*. Boca Raton: CRC Press.
- SCANLON, J. D. & MACKNESS, B. S. (2002). A new giant python from the Pliocene Bluff Downs Local Fauna of northeastern Queensland. *Alcheringa*, **25**, 425–437.
- SCHAAL, S. (2004). *Palaeopython fischeri* n. sp. (Serpentes: Boidae), eine Riesenschlange aus dem Eozän (MP 11) von Messel. *Courier Forschungsinstitut Senckenberg*, **252**, 35–45.
- SCHINZ, H. R. (1833–1835). *Naturgeschichte und Abbildungen der Reptilien. Nach den neuesten Systemen zum gemeinnützigen Gebrauche entworfen und mit Berücksichtigung für den Unterricht der Jugend bearbeitet. Nach der Natur und dem vorzüglichsten Originalien gezeichnet und lithographirt von K. J. Brodtmann. Des Thierreichs dritter Theil*. Shaffhausen, Brodtman Lithographischer Anhalt, 240 pp.
- SCHLEGEL, H. (1837). *Essai sur la physionomie des serpens. Partie Descriptive*. La Haye, J. Kips, J. HZ. & W. P. van Stockum, 606 pp.
- SCHLEGEL, H. (1837–1844). *Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworfen herausgegeben und mit einem erläuternden Texte begleitet*. Düsseldorf, Arnz & Comp, 141 pp.
- SCHLEGEL, H. (1851). Description d'une nouvelle espèce du genre eryx, *Eryx reinhardtii*. *Bijdragen tot de Dierkunde*, **3**, 1–3.
- SCLATER, W. L. (1891). *List of snakes in the Indian Museum*. Calcutta, The Indian Museum, 79 pp.
- SCHNEIDER, J. G. (1801). *Historiae amphibiorum naturalis et literariae. Fasciculus secundus, continens Crocodilos, Scincos, Chamaeauras, Boas, Pseudoboas, Elapes, Angues, Amphisbaenas et Caecilias*. Jenae (= Jena), Friederici Frommann, 364 pp.
- SCHNEIDER, J. G. (1821). Beytrag zur Klassifikation und kritische Uebersicht der Arten aus der Gattung der Riesenschlangen (Boa). *Abhandlung Denkschriften der Königlichen Akademie der Wissenschaften zu München*, **7**, 89–134.
- SCHREIBER, E. (1875). *Herpetologia Europaea : eine systematische Bearbeitung der Amphibien und Reptilien welche bisher in Europa aufgefunden sind*. Braunschweig, F. Vieweg, 639 pp.
- SCHUBERT, G. H. VON (1837). *Die Geschichte der Natur: als zweite gänzlich umgearbeitete Auflage der allgemeinen Naturgeschichte*. Erlangen, Bei J. J. Palm und Ernst Enke, 593 pp.
- SEETZEN, U. J. (1796). Ophiologische Fragmente. *Meyer's Zoologische Archiv*, **2**, 49–74.
- SHAW, G. (1802). *General Zoology, or Systematic Natural History. Vol. III. Part II. Amphibia*. London, G. Kearsley, pp. 313–615.

- SHERBORN, C. D. (1914). An attempt at a fixation of the dates of issue of the parts of the publications of the Musée d'Histoire Naturelle of Paris, 1802–1850. *Annals and Magazine of Natural History, series 8*, **13**, 365–368.
- SIMPSON, G. G. (1933). A new fossil snake from the *Notostylops* Beds of Patagonia. *Bulletin of the American Museum of Natural History*, **67**, 1–22.
- SLOWINSKI, J. B. & LAWSON, R. (2002). Snake phylogeny: evidence from nuclear and mitochondrial genes. *Molecular Phylogenetics and Evolution*, **23**, 194–202.
- SMITH, K. T. (2013). New constraints on the evolution of the snake clades *Ungaliophiinae*, *Loxocemidae* and *Colubridae* (Serpentes), with comments on the fossil history of erycine boids in North America. *Zoologischer Anzeiger*, **252**, 157–182.
- SMITH, K. T., ČERŇANSKÝ, A., SCANFERLA, A. & SCHAALE, S. F. K. (2018). Lizards and snakes – warmth-loving sunbathers, pp. 123–148 in: SMITH, K. T., SCHAALE, S. F. K. & HABERSETZER, J. (eds.) *Messel: An Ancient Greenhouse Ecosystem*. Stuttgart, Schweizerbart.
- SMITH, K. T. & SCANFERLA, A. (2016). Fossil snake preserving three trophic levels and evidence for an ontogenetic dietary shift. *Palaeobiodiversity and Palaeoenvironments*, **96**, 589–599.
- SONNINI, C. S. & LATREILLE, P. A. (1801). *Histoire naturelle des reptiles, avec figures dessinées d'après nature. Tome IV. Seconde partie. Serpens*. Paris, Deterville, 410 pp.
- SPIX, J. DE & WAGLER, J. (1824). *Serpentum brasiliensium species novae ou histoire naturelle des espèces nouvelles de serpens, recueillies et observées pendant le voyage dans l'intérieur du Brésil dans les années 1817, 1818, 1819, 1820, exécuté par ordre de sa majesté le Roi de Bavière*. Monachii (= Munich), Typis F. S. Hübschmanni, 75 pp.
- STANNIUS, H. (1856). *Zoönomie der Amphibien*, 2nd ed. Berlin, Von Veit.
- STREICHER, J. W. & WIENS, J. J. (2016). Phylogenomic analyses reveal novel relationships among snake families. *Molecular Phylogenetics and Evolution*, **100**, 160–169.
- STROMER, E. F. VON (1910). 39. Reptilien- und Fischreste aus dem marinen Alttertiär Südtogo (Westafrika). *Zeitschrift der deutschen geologischen Gesellschaft*, **62**, 478–507.
- STROMER, E. F. VON (1912). *Lehrbuch der paläozoologie. II. Teil: wirbeltiere*. Leipzig & Berlin, B.G. Teubner, 325 pp.
- SWAINSON, W. (1839). On the natural history and classification of fishes, amphibians and reptiles, or monocardian animals. in: *The Cabinet Cyclopædia, Natural History. Volume II*. London, Longman, Orme, Brown, Green & Longmans, 452 pp.
- SZYNDLAR, Z. (1991). A review of Neogene and Quaternary snakes of Central and Eastern Europe. Part I: *Scolecophidia*, *Boidae*, *Colubrinae*. *Estudios Geológicos*, **47**, 103–126.
- SZYNDLAR, Z. & BÖHME, W. (1996). Redescription of *Tropidonotus atavus* von Meyer, 1855 from the upper Oligocene of Rott (Germany) and its allocation to *Rottophis* gen. nov. (Serpentes, Boidae). *Palaeontographica A*, **240**, 145–161.
- SZYNDLAR, Z. & RAGE, J.-C. (2003). *Non-erycine Booidea from the Oligocene and Miocene of Europe*. Kraków, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, 111 pp.
- SZYNDLAR, Z. & SCHLEICH, H.-H. (1993). Description of Miocene snakes from Petersbuch 2 with comments on the lower and middle Miocene ophidian faunas of southern Germany. *Stuttgarter Beiträge zur Naturkunde B*, **192**, 1–47.
- SZYNDLAR, Z. & SCHLEICH, H.-H. (1994). Two species of the genus *Eryx* (Serpentes; Boidae; Erycinae) from the Spanish Neogene with comments on the past distribution of the genus in Europe. *Amphibia-Reptilia*, **15**, 233–248.
- TONINI, J. F. R., BEARD, K. H., FERREIRA, R. B., JETZ, W. & PYRON, R. A. (2016). Fully-sampled phylogenies of squamates reveal evolutionary patterns in threat status. *Biological Conservation*, **204**, 23–31.
- TORRES-CARVAJAL, O., DE QUEIROZ, K. & SCHULTE, J. A. (in press). *Iguanidae* in: DE QUEIROZ, K., CANTINO, P. D. & GAUTHIER, J. A. (eds) *Phylonyms: A Companion to the PhyloCode*. Boca Raton, Florida, CRC Press.
- TROSCHEL, F. H. (1861). Über den Unterkiefer der Schlangen und über die fossile Schlange von Rott. *Archiv für Naturgeschichte*, **27**, 326–360.
- UNDERWOOD, G. (1967). A contribution to the classification of snakes. London, Trustees of the British Museum, **653**, 1–179.
- UNDERWOOD, G. & STIMSON, A. F. (1990). A classification of pythons (Serpentes, Pythoninae). *Journal of Zoology, London*, **221**, 565–603.
- VIDAL, N., DELMAS, A.-S. & HEDGES, B. S. (2007). The higher-level relationships of alethinophidian snakes inferred from seven nuclear and mitochondrial genes, pp. 27–33 in: HENDERSON, R. W. & POWELL, R. (eds.) *Biology of the Boas and Pythons*. Eagle Mountain, Utah, Eagle Mountain Publishing.
- VIDAL, N. & HEDGES, B. S. (2009). The molecular evolutionary tree of lizards, snakes, and amphisbaenians. *Comptes Rendus Biologies*, **332**, 129–139.
- VIDAL, N., RAGE, J.-C., COULOUX, A. & HEDGES, B. S. (2009). Snakes (Serpentes), pp. 390–397 in: Hedges, B. S. & Kumar, S. (eds) *The Timetree of Life*. New York, Oxford University Press.
- WAGLER, J. G. (1828). Auszüge aus seinem Systema Amphibiorum. *Isis von Oken, Leipzig*, **21**, 740–744.
- WALLACH, V. & GÜNTHER, R. (1998). Visceral anatomy of the Malaysian snake genus *Xenophidion*, including a cladistic analysis and allocation to a new family (Serpentes, Xenophidiidae). *Amphibia-Reptilia*, **19**, 385–405.
- WALLACH, V., WILLIAMS, K. L. & BOUNDY, J. (2014). *Snakes of the world: a catalogue of living and extinct species*. Boca Raton, London and New York, CRC Press, 1237 pp.
- WIEGMANN, A. F. A. (1832). *Handbuch der Zoologie*. Berlin, C.G. Lüderitz, 621 pp.
- WIENS, J. J., HUTTER, C. R., MULCAHY, D. G., NOONAN, B. P., TOWNSEND, T. M., SITES, J. W. & REEDER, T. W. (2012). Resolving the phylogeny of lizards and snakes (Squamata) with extensive sampling of genes and species. *Biology Letters*, **8**, 1043.
- WILCOX, T. P., ZWICKL, D. J., HEATH, T. A. & HILLIS, D. M. (2002). Phylogenetic relationships of the dwarf boas and a comparison of Bayesian and bootstrap measures of phylogenetic support. *Molecular Phylogenetics and Evolution*, **25**, 361–371.
- ZACHARIAS, H. C. E. (1898). Die Phylogene der Kopfschilder bei den Boiden. *Zoologische Jahrbücher*, **10**, 56–90.
- ZAHER, H. (1994). Les Tropidopoeoidea (Serpentes; Alethinophidia) sont-ils réellement monophylétiques? Arguments en faveur de leur polyphylétisme. *Comptes rendus de l'Académie des Sciences, Paris*, **317**, 471–478.
- ZAHER, H., GRAZZIOTIN, F. G., CADLE, J. E., MURPHY, R. W., CESAR DE MOURA-LEITE, J. & BONATTO, S. L. (2009). Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de Zoologia*, **49**, 115–153.
- ZAHER, H., MURPHY, R. W., ARREDONDO, J. C., GRABOSKI, R., MACHADO-FILHO, P. R., MAHLOW, K., MONTINGELLI, G. G., BOT-TALLO QUADROS, A., ORLOV, N. L., WILKINSON, M., ZHANG, Y.-P. & GRAZZIOTIN, F. G. (2019). Large-scale molecular phylogeny, morphology, divergence-time estimation, and the fossil record of advanced caenophidian snakes (Squamata: Serpentes). *PLoS ONE*, **14**, e0216148.
- ZAHER, H. & SCANFERLA, A. C. (2012). The skull of the Upper Cretaceous snake *Dinilysia patagonica* Smith-Woodward, 1901, and its phylogenetic position revisited. *Zoological Journal of the Linnean Society*, **164**, 194–238.
- ZENNECK, J. (1898). Die Zeichnung der Boiden. *Zeitschrift für Wissenschaftliche Zoologie, Leipzig*, **64**, 1–384.
- ZHENG, Y. & WIENS, J. J. (2016). Combining phylogenomic and supermatrix approaches, and a time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162 species. *Molecular Phylogenetics and Evolution*, **94**, 537–547.
- ZITTEL, K. A. (1887–1890). *Handbuch der Paläontologie. Palaeozoologie. III. Pisces, Amphibia, Reptilia, Aves*. München and Leipzig, Druck und Verlag von R. Oldenbourg, 900 pp.